

**Amendments to the Claims**

This listing of claims will replace all prior versions, and listings of claims in the application:

**Listing of Claims:**

1 (currently amended). A polynucleotide comprising a sequence encoding a polypeptide that is capable of high level presentation of antigenic peptides on antigen-presenting cells, wherein the polypeptide comprises a  $\beta$ 2-microglobulin molecule that is linked through its carboxyl terminal to a polypeptide stretch that allows the anchorage of the  $\beta$ 2-microglobulin molecule to the cell membrane selected from the group consisting of the human CD3  $\zeta$  polypeptide, the MHC I class heavy chain of HLA-A, HLA-B or HLA-C molecule and CD40, and through its amino terminal to at least one antigenic peptide comprising an MHC class I epitope, wherein said antigenic peptide is not ~~related to an autoimmune disease~~ capable of activating autoreactive cells causing an autoimmune disease.

2 (original). The polynucleotide of claim 1, wherein said polypeptide stretch at the  $\beta$ 2-microglobulin carboxyl terminal consists of a bridge peptide which spans the whole distance to the cell membrane, said bridge peptide being linked to a sequence which can exert the required anchoring function.

3(original). The polynucleotide of claim 2, wherein said bridge peptide is the peptide of SEQ ID NO: 1, of the sequence: LRWEPSSQPTIPI.

Claim 4 (cancelled).

5(currently amended). The polynucleotide of claim 4 1, wherein said bridge peptide is linked to the transmembrane and cytoplasmic domains from the MHC class I heavy chain HLA-A2 molecule, of the SEQ ID NO: 2, of the sequence:  
VGIIAGLVLFGLAVITGAVVAAVMWRRKSSDRKGGSYSQAASSDSAQGSVDVSLTACKV

Claims 6-10 (cancelled).

11(previously presented). The polynucleotide of claim 1, wherein said at least one antigenic peptide comprising a MHC class I epitope is linked to the  $\beta$ 2-microglobulin amino terminal through a peptide linker.

12(original). The polynucleotide of claim 1, wherein said at least one antigenic peptide is at least one antigenic determinant of one sole antigen.

13(original). The polynucleotide of claim 11, wherein said at least one antigenic peptide is at least one antigenic determinant of each one of at least two different antigens.

14(previously presented). The polynucleotide of claim 12, wherein said antigen is a tumor-associated antigen (TAA).

15(original). The polynucleotide of claim 14, wherein said TAA is selected from the group consisting of alpha-fetoprotein, BA-46/lactadherin, BAGE, BCR-ABL fusion protein, beta-catenin, CASP-8, CDK4, CEA, CRIPTO-1, elongation factor 2, ETV6-AML1 fusion protein, G250, GAGE, gp100, HER-2/neu, intestinal carboxyl esterase, KIAA0205, MAGE, MART-1/Melan-A, MUC-1, N-ras, p53, PAP, PSA, PSMA, telomerase, TRP-1/gp75, TRP-2, tyrosinase, and uroplakin Ia, Ib, II and III.

16(previously presented). The polynucleotide of claim 15, wherein said antigenic peptide is selected from the group consisting of:

- (i) the alpha-fetoprotein peptide  
GVALQTMKQ (SEQ ID NO:4);
- (ii) the BAGE-1 peptide AARAVFLAL (SEQ ID  
NO:5);
- (iii) the BCR-ABL fusion protein peptide  
SSKALQRPV (SEQ ID NO:6);
- (iv) the beta-catenin peptide SYLDSGIHF  
(SEQ ID NO:7);
- (v) the CDK4 peptide ACDPHSGHFV (SEQ ID  
NO:8);

- (vi) the CEA peptide YLSGANLNL (SEQ ID NO:9);
- (vii) the elongation factor 2 peptide ETVSEQSNV (SEQ ID NO:10);
- (viii) the ETV6-AML1 fusion protein peptide RIAECILGM (SEQ ID NO:11);
- (ix) the G250 peptide HLSTAFARV (SEQ ID NO:12);
- (x) the GAGE-1,2,8 peptide YRPRPRRY (SEQ ID NO:13);
- (xi) the gp100 peptides KTWGQYWQV (SEQ ID NO:14), A)MLGTHTMEV (SEQ ID NO:15), ITDQVPFSV (SEQ ID NO:16), YLEPGPVTA (SEQ ID NO:17), LLDGTATLRL (SEQ ID NO:18), VLYRYGSFSV (SEQ ID NO:19), SLADTNSLAV (SEQ ID NO:20), RLMKQDFS (SEQ ID NO:21), RLPRIFCSC (SEQ ID NO:22), LIYRRRLMK (SEQ ID NO:23), ALLAVGATK (SEQ ID NO:24), IALNFPQSQK (SEQ ID NO:25) and ALNFPQSQK (SEQ ID NO:26);
- (xii) the HER-2/neu peptide KIFGSLAFL (SEQ ID NO:27);

- (xiii) the intestinal carboxyl esterase  
peptide SPRWWPTCL (SEQ ID NO:28);
- (xiv) the KIAA0205 peptide AEPINIQTW (SEQ  
ID NO:29);
- (xv) the MAGE-1 peptides EADPTGHSY (SEQ ID  
NO:30) and SLFRAVITK (SEQ ID NO:31);
- (xvi) the MAGE-3 peptides EVDPIGHLY (SEQ ID  
NO:32) and FLWGPRALV (SEQ ID NO:33);
- (xvii) the MART-1/Melan-A peptide  
(E)AAGIGILTV (SEQ ID NO:34);
- (xviii) the MUC-1 peptide STAPPVHNV (SEQ ID  
NO:35);
- (xix) the N-ras peptide ILDTAGREEY (SEQ ID  
NO:36);
- (xx) the p53 peptide LLGRNSFEV (SEQ ID  
NO:37);
- (xxi) the PSA peptides FLTPKKLQCV (SEQ ID  
NO:38) and VISNDVCAQV (SEQ ID NO:39);
- (xxii) the telomerase peptide ILAKFLHWL (SEQ  
ID NO:40);
- (xxiii) the TRP-1 peptide MSLQRQFLR (SEQ ID  
NO:41);

- (xxiv) the TRP-2 peptides LLGPGRPYR (SEQ ID NO:42), SVYDFFVWL (SEQ ID NO:43), and TLDSQVMSL (SEQ ID NO:44);
- (xxv) the TRP2-INT2 peptide EVISCKLIKR (SEQ ID NO:45); and
- (xxvi) the tyrosinase peptide KCDICTDEY (SEQ ID NO:46).

17(previously presented). The polynucleotide of claim 14, wherein said at least one antigenic peptide is at least one antigenic determinant of one sole tumor-associated antigen.

18(original). The polynucleotide of claim 17, wherein said at least one antigenic peptide is at least one HLA-A2 binding peptide and at least one HLA-A3 binding peptide derived from the melanoma-associated antigen gp100.

19(original). The polynucleotide of claim 18, wherein said at least one HLA-A2 binding peptide derived from gp100 is selected from the group consisting of SEQ ID NO: 14, 15, 16, 17, 18, 19, 20, 21 and 22, and said at least one gp100 HLA-A3 binding peptide is selected from the group consisting of SEQ ID NO: 23, 24, 25 and 26.

20(previously presented). The polynucleotide of claim 14, wherein said at least one antigenic peptide is at least one

antigenic determinant of each one of at least two different tumor-associated antigens.

21(original). The polynucleotide of claim 20, wherein said at least one antigenic peptide is at least one HLA-A2 binding peptide derived from each one of the melanoma associated antigens gp100 and Melan-A/MART-1.

22(original). The polynucleotide of claim 21, wherein said at least one antigenic peptide is at least one HLA-A3-restricted gp100 and at least one HLA-A2-restricted Melan-A/MART-1 peptide.

23(previously presented). The polynucleotide of claim 12, wherein said antigen is an antigen from a pathogen selected from the group consisting of a bacterial, viral, fungal and parasite antigen.

24(original). The polynucleotide of claim 23 wherein the antigen is a viral antigen.

25(original). The polynucleotide of claim 24 wherein the viral antigen is an HIV protein selected from the group consisting of the HIV-1 regulatory proteins Tat and Rev and the HIV envelope protein, in which case the antigenic peptide derived therefrom has the sequence RGPGRAFVTI (SEQ ID NO:47).

26(original). The polynucleotide of claim 11, wherein said at least one antigenic peptide is at least one idiotypic peptide expressed by autoreactive T lymphocytes.

27(original). The polynucleotide of claim 26, wherein said at least one idiotypic peptide is derived from a CDR (complementarity-determining region) sequence of an immunoglobulin or of a TCR chain, optionally containing said CDR flanking segments.

28(original). The polynucleotide of claim 27, wherein said CDR is the CDR3 of an immunoglobulin or of a TCR chain.

29. (previously presented) The polynucleotide of claim 1 that is an expression vector.

30(previously presented). An expression vector comprising a polynucleotide according to claim 1.

31(original). A recombinant viral vector of claim 30.

32(currently amended). An antigen-presenting cell transfected with a polynucleotide comprising a sequence encoding a polypeptide comprising a  $\beta$ 2-microglobulin molecule that is linked through its carboxyl terminal to a polypeptide stretch that allows the anchorage of the  $\beta$ 2-microglobulin molecule to the cell membrane selected from the group consisting of the human CD3



ζ polypeptide, the MHC I class heavy chain of HLA-A, HLA-B or HLA-C molecule and CD40, and through its amino terminal to at least one antigenic peptide comprising a MHC class I epitope.

33(original). The antigen-presenting cell of claim 32 selected from the group consisting of a dendritic cell, a macrophage, a B cell and a fibroblast.

34(previously presented). The antigen-presenting cell of claim 32 wherein said antigenic peptide is a peptide not related to an autoimmune disease.

35(original). The antigen-presenting cell of claim 34, wherein said antigenic peptide is at least one peptide derived from at least one TAA.

36(original). The antigen-presenting cell of claim 34, wherein said antigenic peptide is at least one peptide derived from an antigen from a pathogen selected from the group consisting of a bacterial, a viral, a fungal and a parasite antigen.

37(previously presented). A DNA vaccine comprising a polynucleotide of claim 1 or an expression vector of claim 30.

38(currently amended). The DNA vaccine of claim 37 for prevention or treatment of cancer wherein said polynucleotide is

a polynucleotide comprising a sequence encoding a polypeptide that is capable of high level presentation of antigenic peptides on antigen-presenting cells, wherein the polypeptide comprises a  $\beta$ 2-microglobulin molecule that is linked through its carboxyl terminal to a polypeptide stretch that allows the anchorage of the  $\beta$ 2-microglobulin molecule to the cell membrane selected from the group consisting of the human CD3  $\zeta$  polypeptide, the MHC I class heavy chain of HLA-A, HLA-B or HLA-C molecule and CD40, and through its amino terminal to at least one antigenic peptide comprising a MHC class I epitope, and said at least one antigenic peptide is at least one antigenic determinant of one sole tumor-associated antigen (TAA).

39(currently amended). The DNA vaccine of claim 37 for prevention or treatment of a disease caused by a pathogenic organism wherein said polynucleotide is a polynucleotide comprising a sequence encoding a polypeptide that is capable of high level presentation of antigenic peptides on antigen-presenting cells, wherein the polypeptide comprises a  $\beta$ 2-microglobulin molecule that is linked through its carboxyl terminal to a polypeptide stretch that allows the anchorage of the  $\beta$ 2-microglobulin molecule to the cell membrane selected from the group consisting of the human CD3  $\zeta$  polypeptide, the MHC I class heavy chain of HLA-A, HLA-B or HLA-C molecule and CD40, and

through its amino terminal to at least one antigenic peptide comprising a MHC class I epitope, and said at least one antigenic peptide is at least one antigenic determinant of one sole antigen from a pathogen selected from the group consisting of a bacterial, viral, fungal and parasite antigen.

40(original). A cellular vaccine, which comprises an antigen presenting cell of claim 32.

41(previously presented). The cellular vaccine of claim 40 wherein the antigen presenting cell is selected from the group consisting of a dendritic cell, a macrophage, a B cell and a fibroblast.

42(currently amended). The cellular vaccine of claim 41, wherein the at least one antigenic peptide presented by the antigen presenting cell is a peptide not ~~related to an autoimmune disease~~ capable of activating autoreactive cells causing an autoimmune disease.

43(original). The cellular vaccine of claim 42 for prevention or treatment of cancer wherein the antigen presenting cell presents at least one peptide derived from at least one tumor associated antigen.

44(original). The cellular vaccine of claim 42 for prevention or treatment of a disease caused by a pathogenic

organism wherein the antigen presenting cell presents at least one peptide derived from a pathogenic antigen.

Claims 45-46 (cancelled).

47(currently amended). A method of immunizing a mammal against a tumor-associated antigen comprising the step of immunizing the mammal with a cellular vaccine, which comprises an antigen presenting cell transfected with a polynucleotide comprising a sequence encoding a polypeptide comprising a  $\beta$ 2-microglobulin molecule that is linked through its carboxyl terminal to a polypeptide stretch that allows the anchorage of the  $\beta$ 2-microglobulin molecule to the cell membrane selected from the group consisting of the human CD3  $\zeta$  polypeptide, the MHC I class heavy chain of HLA-A, HLA-B or HLA-C molecule and CD40, and through its amino terminal to at least one antigenic peptide comprising a MHC class I epitope, wherein said antigen presenting cell is selected from the group consisting of a dendritic cell, a macrophage, a B cell or a fibroblast, and said at least one antigenic peptide is at least one peptide derived from at least one tumor-associated antigen.

48(currently amended). A method of immunizing a mammal against a disease caused by a pathogenic organism comprising the step of immunizing the mammal with a cellular vaccine, which comprises an antigen presenting cell transfected

with a polynucleotide comprising a sequence encoding a polypeptide comprising a  $\beta$ 2-microglobulin molecule that is linked through its carboxyl terminal to a polypeptide stretch that allows the anchorage of the  $\beta$ 2-microglobulin molecule to the cell membrane selected from the group consisting of the human CD3  $\zeta$  polypeptide, the MHC I class heavy chain of HLA-A, HLA-B or HLA-C molecule and CD40, and through its amino terminal to at least one antigenic peptide comprising a MHC class I epitope, wherein said antigen presenting cell is selected from the group consisting of a dendritic cell, a macrophage, a B cell or a fibroblast, and said at least one antigenic peptide is at least one antigenic peptide derived from a pathogen.

49(previously presented). A pharmaceutical composition comprising as an active ingredient at least one polynucleotide of claim 1 and a pharmaceutically acceptable carrier.

50(original). The pharmaceutical composition of claim 49 wherein the polynucleotide comprises a sequence encoding a polypeptide comprising at least one antigenic peptide derived from at least one tumor associated antigen.

51(original). The pharmaceutical composition of claim 49 wherein the polynucleotide comprises a sequence encoding a

Appln. No. 10/517,784  
Amd. dated February 12, 2008  
Reply to Office Action of February 7, 2008

polypeptide comprising at least one antigenic peptide derived from a pathogenic antigen.

52 (previously presented). A pharmaceutical composition comprising as an active ingredient at least one antigen presenting cell of claim 32 and a pharmaceutically acceptable carrier.

53 (previously presented). The polynucleotide of claim 13, wherein said antigen is a tumor-associated antigen (TAA).

Claims 54 and 55 (cancelled).